

Impact of genomic selection on genetic diversity in 5 European local cattle breeds

EAAP 2023

Renzo Bonifazi, Gwendal Restoux, Pascal Croiseau, Theo Meuwissen, Stephanie Minery, Jack Windig



Acknowledgments



- Breeding companies for providing pedigree and SNP data
- Gerben de Jong (CRV)
- Mathijs van Pelt (CRV)



TOWARDS IMPROVEMENT OF **RUMINANT BREEDING**
THROUGH **GENOMIC AND EPIGENOMIC APPROACHES**

- This project has received funding from the European Union's Horizon 2020 Programme for Research & Innovation under grant agreement n°101000226.
- This project is part of EuroFAANG (<https://eurofaang.eu>).


Genetic diversity in local cattle populations

- Strong selection can lead to high inbreeding rates
 - associated los of genetic diversity
 - + inbreeding depression
 - + risk expression genetic defect
- Especially in small populations
- Genomic selection (GS) now applied in local breeds
- Main question
 - Did inbreeding rates change after the introduction of GS in 5 local breeds?



Changes in inbreeding rate after introduction GS

- Genomic selection (GS): starting ~2008 (VanRaden, 2008)
- Larger genetic gains, shorter generation intervals
- Expectation ΔF to go down



Up

- Holstein

- the Netherlands (Doekes et al. 2018)
- USA (Forutan et al. 2018; Makanjuola et al. 2020)
- Poland (Topolski and Jagusiak 2020)
- France (Doublet et al. 2019)
- Australia (Scott et al. 2021)
- Italy (Ablondi 2022)

- Jersey (Makanjuola et al. 2020)

- Finnish Ayrshire (Sarviaaho et al. 2023)

Not up

- Normande

- Mont Beliarde

- Doublet et al. 2019

- Aberdeen Angus

- Lozada-Soto et al. 2021

5 cattle populations



MRY (NL)

Abondance (F)



Tarentaise (F)



Norwegian Red (N)

Vosgienne (F)




Some data

Breed	Calf/year	Pedigree	Genotypes	Years analysed	Introduction GS
MRIJ	5,053	205,934	4,645	1975-2021	2018
Norwegian Red	35,701	721,805	193,489	2000-2020	2016
Abondance	25,020	807,387	16,427	2000-2020	2014
Tarentaise	9,514	259,150	8,882	2000-2020	2014
Vosgienne	3,738	99,833	4,466	2000-2020	2014

MRY inbreeding and kinship levels


- All animals in pedigree



- Steady increase up to 2013
- Decrease 2013 – 2018
- Increase since introduction GS in 2019


MRY inbreeding and kinship levels

- Genotyped animals only



Norwegian Red inbreeding and kinship levels


- All animals in pedigree



- Steady increase of inbreeding
- Kinship decreases since 2009
- Inbreeding continues to increase since introduction GS in 2016
- Kinship continues to decrease

Norwegian Red inbreeding and kinship levels


- Genotyped animals only



- F(ROH) stable, slight increase since introduction GS
- Kinship decrease after 2014, stable after 2016
- F(ped) increase but less after 2014

French breeds

- Steady increase in all levels
- After introduction GS (2014)
 - Abondance
 - Less steep
 - Especially F(ROH)
 - Tarentaise
 - Steeper
 - especially F(Ped)
 - Vosgienne
 - Clearly less steep
 - Especially F(ROH)




Change in inbreeding and kinship rates

Breed	F_{ped}	F_{ROH}	f
MRIJ all	Stable/Up	-	Stable/Up
MRIJ genotyped	UP	UP	Stable/Up
NR all	Stable	-	Down/up
NR genotyped	Down/Stable	UP	Down/up
Abondance	Down	Down	Down
Tarentaise	Up	Up	Up
Vosgienne	Down	Down	Down

- Mixed pattern: rates can go
 - up
 - or down
 - or remain stable
- after introduction of GS

Genetic management can reduce kinship



- MRIJ and Norwegian Red both use optimal contributions to select sires
- After introduction of OC a clear reduction in kinship

Summary

- Inbreeding rates (ΔF_{gen}):
 - Increased (MRIJ, Tarentaise)
 - More or less stable (Norwegian Red)
 - Decreased (Abondance, Vosgienne)
- Genomic Selection:
 - shorter generation intervals
 - screening more individuals
 - preselection of less sires

Main take-home message

- **genetic management** of local breeds more important than implementation of genomic selection per se

