INTRODUCTION

The latest genomic routine international evaluation for **females fertility** traits took place as scheduled at the Interbull Centre. Data from 18 countries were included in this evaluation.

International genetic evaluations for female fertility traits of bulls from Australia, Austria, Belgium, Canada, Czech Republic, Denmark, Finland, France, Germany, Ireland, Israel, Italy, Netherlands, New Zealand, Norway, Poland, Spain, Sweden, Switzerland, South Africa, the United Kingdom and the United States of America were computed. Holstein data were included in this evaluation.

BEL, CAN, DEU, ESP, FRA, DFS, GBR, ITA, NLD, POL submitted GEBVs.

cc1:CAN, DEU,FRA, DFS, GBR, ITA, NLD, POLcc2:BEL, CAN, DEU, ESP, FRA, DFS, GBR, ITA, NLD, POLcrc:BEL, CAN, DEU, ESP, FRA, DFS, GBR, ITA, NLD, POLhco:CAN, DEU,FRA, DFS,int:BEL, CAN, DEU, ESP,NLD, POL

Based on a decision made by Interbull Steering committee in August 2007, female fertility traits are classified as follows:

- T1 (HC): Maiden (H)eifer's ability to (C)onceive. A measure of confirmed conception, such as conception rate (CR), will be considered for this trait group. In the absence of confirmed conception an alternative measure, such as interval first-last insemination (FL), interval first insemination-conception (FC), number of inseminations (NI), or non-return rate (NR,preferably NR56) can be submitted;
- T2 (CR): Lactating (C)ow's ability to (R)ecycle after calving. The interval calving-first insemination (CF) is an example for this ability. In the abscence of such a trait, a measure of the interval calving-conception, such as says oprn (DO) or calving interval (CI) can be submitted;
- T3 (C1): Lactating (C)ow's ability to conceive (1), expressed as a rate trait. Traits like conception rate (CR) and non-return rate (NR, preferably NR56) will be considered for this trait group;
- T4 (C2): Lactating (C)ow's ability to conceive (2), expressed as an interval trait. The interval first insemination-conception (FC) or interval first-last insemination (FL) will be considered for this trait group. As an alternative, number of inseminations (NI) can be submitted. In the abscence of any of these traits, a measure of interval calving-conception such as days open (DO), or calving interval (CI) can be submitted. All countries are expected to submit data for this trait group, and as a last resort the trait submitted under T3 can be submitted for T4 as well.
- T5 (IT): Lactating cow's measurements of (I)nterval (T)raits calving-conception, such as days open (DO) and calving interval (CI).

Based on the above trait definitions the following traits have been submitted for international genetic evaluation of female fertility traits.

Country Traits Submitted traits and their definitions

AUS	T2=CY T4=C2 T5=IT	Calving interval converted to 42 days pregnancy rate Calving interval converted to 42 days pregnancy rate Calving interval converted to 42 days pregnancy rate
BEL	T2=CY T4=C2 T5=IT	PR=Pregnancy Rate (=[21/(DO-45+11)]*100, with DO=days open) PR=Pregnancy Rate (=[21/(DO-45+11)]*100, with DO=days open) PR=Pregnancy Rate (=[21/(DO-45+11)]*100, with DO=days open)
CAN	T1=HC T2=CY T3=C1 T4=C2 T5=IT	NR=Non Return Rate after 56 Days in heifers (NRR), % CF=Interval from Calving to First Service in cows(CF) NR=Non Return Rate after 56 Days in cows(NRR), % FC=Interval first insemination-conception in cows DO=Days open

- CR=Heifers' Conception rate T1=HC CHE T2=CR CF=Interval from Calving to First Service (ICF), days T3=C1 NR=Non Return Rate after 56 Days (NRR), % T4=C2 NR=Non Return Rate after 56 Days (NRR), % CZE T1=HC CR=Heifers' Conception rate (pregnant or not after 3 months) T3=C1 CR=Cows' Conception rate (pregnant or not after 3 months) T4=C2 CR=Cows' Conception rate (pregnant or not after 3 months) AUT/DEU T1=HC NR=Heifers' Non Return Rate after 56 days T2=CY CF=Interval from calving to first insemination cows (days) T3=C1 NR=Cows' Non Return Rate after 56 days T4=C2 FL=Interval from first to last insemination cows (days) DO=Days open (days) T5=IT T1=HC NR=Heifers' Non Return Rate after 56 days DFS T2=CY CF=Interval from calving to first insemination cows (days) T3=C1 NR=Cows' Non Return Rate after 56 days T4=C2 FL=Interval from first to last insemination cows (days) T5=IT DO=Days open (days) T2=CY DO=Days open ESP T4=C2 DO=Days open T5=IT DO=Days open FRA T1=HC CR=Heifers' Conception rate (binary trait) for maiden heifers T2=CY Interval between calving and first AI T3=C1 CR=Cows' Conception rate (binary trait) for cows T4=C2 FL=Interval from first to last insemination cows (days) T2=CY CI=days between 1st and 2nd calvings GBR T3=C1 NR=1st lactation non return at 56 days T4=C2 CI=days between 1st and 2nd calvings T5=IT CI=days between 1st and 2nd calvings T2=CY CI=Calving interval IRL T4=C2 CI=Calving interval T5=IT CI=Calving interval ISR T3=C1 CR=Inverse of the number of insemination to conception (%) T4=C2 CR=Inverse of the number of insemination to conception (%) T2=CY CF=Days to first service ITA NR=Non-return rate at 56 days (%) T3=C1 T4=C2 CI=Calving Interval (days) T5=IT CI=Calving interval (days) ITA(BSW) T2=CY CF=Interval calving to first insemination T4=C2 Days Open T5=IT CI=Calving interval T1=HC CR=Heifers' Conception rate NLD T2=CY CF=Interval calving to first insemination (days) T3=C1 CR=Cows' Conception rate (binary trait) for cows T4=C2 FL=Interval from first to last insemination cows (days) T5=IT CI=Calving Interval (days) NOR T1=HC NR=NR=Non-return rate 56 days (heifers) T2=CY CF=Interval calving to first insemination (days) T3=C1 NR=NR=Non-return rate 56 days (cows) T4=C2 CI=Calving Interval (days) T5=IT CI=Calving Interval (days) NZL T2=CY PM=Lactating cow's ability to start cycling T4=C2 PC=Lactating cow's ability to conceive (CR42) T5=IT PC=Lactating cow's ability to conceive (CR42)
 - POL T1=HC Non return rate at 56 days for heifer T2=CR Interval from calving to first insemination T3=C1 Non return rate at 56 days for cows

	T4=IT T5=IT	1 1
USA	T1=HC T2=CY T3=C1 T4=C2 T5=IT	CF=Interval from calving to first insemination CR=Conception rate (cow) DP=Daughter Pregnancy Rate
ZAF		CI=Calving Interval CI=Calving Interval
CHANCE		CONAL PROCEDURES
Change	s in the	national genetic evaluation of fertility traits are as follows:
INTERB		SES COMPARED TO THE DECEMBER ROUTINE RUN
No cha	nges in I	Interbull procedures
		OF ANALYSIS
classi GEBVs countr a GMAC From t classi with a genomi	cal EBVs from the ies parti E EBV but hose elev cal MACE further c bulls w	A populations sent GEBV data for up to 38 traits, while for the same traits were used in the analyses. Young bull GEBV providers have been converted to the scales of all cipating in classical MACE. A bull will get a MACE EBV or t not both. Yen countries, National GEBVs of bulls less than seven years of age and with no proofs were included for the breeding value prediction requirement of either a MACE-PA or a GMACE-PA (for young with young genomic sires) being available.
-		
	IFIC LITE	RATURE
		aal genetic evaluation procedure is based on international work ne following scientific publications:
Sulliv	an, P.G.	2016. Defining a Parameter Space for GMACE. Interbull Bulletin 50, p 85-93.
	en, P.M. el. Evol.	and Sullivan, P.G. 2010. International genomic evaluation methods for dairy cattle. 42:7
	an, P.G. in 45, Ar	and Jakobsen, J.H. 2012. Robust GMACE for young bulls methodology. Interbull sticle 1.
	•	2012a. GMACE reliability approximation. Report to the GMACE working group of CE_rels 2013 $$
	an, P.G. vce 2013	2012b. GMACE variance estimation. Report to the GMACE working group of Interbull.
	an, P.G. gedcs 201	2012c. GMACE Weighting Factors. Report to the GMACE working group of Interbull. .3
		and Sullivan, P.G. 2013. Trait specific computation of shared reference population. .ng Nov 2013

NEXT ROUTINE INTERNATIONAL EVALUATION

Dates for next routine run can be found on http://www.interbull.org/ib/servicecalendar

NEXT TEST INTERNATIONAL EVALUATION

Dates for next routine run can be found on http://www.interbull.org/ib/servicecalendar

PUBLICATION OF INTERBULL ROUTINE RUN

Results were distributed by the Interbull Centre to designated representatives in each country. The international evaluation file comprised international proofs expressed on the base and unit of each country included in the analysis. Such records readily provide more information on bull performance in various countries, thereby minimising the need to resort to conversions.

At the same time, all recipients of Interbull results are expected to honour the agreed code of practice, decided by the Interbull Steering Committee, and only publish international evaluations on their own country scale. Evaluations expressed on another country scale are confidential and may only be used internally for research and review purposes.

Table 1. National evaluation dates in GMACE run December 2017

Country	y Date
BEL	20171201
CAN	
DEU	20171205
DFS	20171107
ESP	20171114
GBR	20171027
ITA	
NLD	
	20171110
	20171206
Table 2	2.
Number	of bulls in reference population for hco
CAN 22	 179 N
	748.0 32963.0
	483.0 30627.0 31755.0
	716.0 29660.0 29390.0 31678.0
	848.0 26359.0 26419.0 25604.0 28472.0
NLD 2	554.0 30923.0 30390.0 29483.0 26350.0 31889.0
	of bulls in reference population for crc
BEL 2	
	248.0 29139.0
DEU	984.0 2958.0 34985.0
DFS	891.0 2588.0 32499.0 33600.0
ESP	959.0 2650.0 32965.0 32874.0 33970.0
GBR	867.0 25998.0 2760.0 2447.0 2487.0 26492.0
ITA 1	089.0 25149.0 1989.0 1589.0 1630.0 23948.0 25553.0
NLD	973.0 2699.0 32869.0 32361.0 32934.0 2500.0 1822.0 34422.0
POL 1	115.0 2814.0 28320.0 28324.0 28777.0 2361.0 1881.0 28406.0 29753.0
FRA 1	004.0 2888.0 31382.0 31054.0 31638.0 2686.0 1863.0 31267.0 27313.0 33395.

_____ Number of bulls in reference population for cc1 _____ CAN 29144.0 DEU 2933.0 33240.0 DFS 2545.0 30757.0 31780.0 FRA 2861.0 29849.0 29466.0 31781.0 GBR 25840.0 2754.0 2429.0 2671.0 26323.0 ITA 25129.0 1975.0 1575.0 1853.0 23888.0 25532.0 NLD 2670.0 31103.0 30542.0 29663.0 2480.0 1806.0 32192.0 POL 2927.0 26567.0 26518.0 25754.0 2378.0 1914.0 26587.0 28561.0 _____ Number of bulls in reference population for cc2 _____ BEL 2607.0 CAN 1395.0 31302.0 DEU 1003.0 2942.0 34950.0 DFS 900.0 2575.0 32465.0 33468.0 ESP 970.0 2635.0 32943.0 32771.0 33870.0 GBR 911.0 28004.0 2714.0 2395.0 2434.0 28476.0 ITA 1147.0 26842.0 1989.0 1593.0 1634.0 25600.0 27240.0 NLD 999.0 2735.0 32898.0 32371.0 32951.0 2506.0 1840.0 34628.0 POL 1479.0 2942.0 28298.0 28207.0 28684.0 2334.0 1933.0 28437.0 30266.0 _____ Number of bulls in reference population for int. _____ BEL 1739.0 848.0 29052.0 CAN DEU 934.0 2461.0 28795.0 DFS 850.0 2060.0 26489.0 27309.0 ESP 914.0 2114.0 26897.0 26638.0 27628.0 GBR 845.0 27465.0 2341.0 1976.0 2012.0 27922.0 ITA 750.0 25543.0 1845.0 1468.0 1506.0 25416.0 25937.0 NLD 932.0 2218.0 26900.0 26318.0 26830.0 2101.0 1706.0 28442.0 POL 990.0 2019.0 22343.0 22143.0 22589.0 1910.0 1491.0 22372.0 23231.0