## INTRODUCTION

\_\_\_\_\_

The latest genomic test international evaluation for females fertility traits took place as scheduled at the Interbull Centre. Data from 20 countries were included in this evaluation.

International genetic evaluations for fertility traits of bulls were computed from: AUS BEL CAN CHE CZE DEU DFS ESP FRA GBR IRL ISR ITA NLD NZL POL USA ZAF URY JPN Holstein data were included in this evaluation.

BEL, CAN, DEU, FRA, DFS, GBR, ITA, NLD, POL submitted GEBVs.

cc1:CAN, DEU,FRA,DFS,GBR,ITA,NLD,POLcc2:BEL,CAN,DEU,ESP,,DFS,GBR,ITA,NLD,POLcrc:BEL,CAN,DEU,ESP,FRA,DFS,GBR,ITA,NLD,POLhco:CAN,DEU,,FRA,DFS,,ITA,NLD,POLint:BEL,CAN,DEU,,FRA,DFS,,ITA,NLD,POL

Based on a decision made by Interbull Steering committee in August 2007, female fertility traits are classified as follows:

T1 (HC): Maiden (H)eifer's ability to (C)onceive. A measure of confirmed conception, such as conception rate (CR), will be considered for this trait group. In the absence of confirmed conception an alternative measure, such as interval first-last insemination (FL), interval first insemination-conception (FC), number of inseminations (NI), or non-return rate (NR,preferably NR56) can be submitted;

- T2 (CR): Lactating (C)ow's ability to (R)ecycle after calving. The interval calving-first insemination (CF) is an example for this ability. In the abscence of such a trait, a measure of the interval calving-conception, such as says oprn (DO) or calving interval (CI) can be submitted;
- T3 (C1): Lactating (C)ow's ability to conceive (1), expressed as a rate trait. Traits like conception rate (CR) and non-return rate (NR, preferably NR56) will be considered for this trait group;
- T4 (C2): Lactating (C) ow's ability to conceive (2), expressed as an interval trait. The interval first insemination-conception (FC) or interval first-last insemination (FL) will be considered for this trait group. As an alternative, number of inseminations (NI) can be submitted. In the abscence of any of these traits, a measure of interval calving-conception such as days open (DO), or calving interval (CI) can be submitted. All countries are expected to submit data for this trait group, and as a last resort the trait submitted under T3 can be submitted for T4 as well.
- T5 (IT): Lactating cow's measurements of (I)nterval (T)raits calving-conception, such as days open (DO) and calving interval (CI).

Based on the above trait definitions the following traits have been submitted for international genetic evaluation of female fertility traits.

\_\_\_\_\_

Country Traits Submitted traits and their definitions \_\_\_\_\_ T2=CY Calving interval converted to 42 days pregnancy rate AUS T4=C2 Calving interval converted to 42 days pregnancy rate T5=IT Calving interval converted to 42 days pregnancy rate T2=CY PR=Pregnancy Rate (=[21/(D0-45+11)]\*100, with D0=days open) BEL T4=C2 PR=Pregnancy Rate (=[21/(DO-45+11)]\*100, with DO=days open) T5=IT PR=Pregnancy Rate (=[21/(DO-45+11)]\*100, with DO=days open) CAN T1=HC NR=Non Return Rate after 56 Days in heifers (NRR), % T2=CY CF=Interval from Calving to First Service in cows(CF) T3=C1 NR=Non Return Rate after 56 Days in cows(NRR), % T4=C2 FC=Interval first insemination-conception in cows T5=IT DO=Days open

- T1=HC CR=Heifers' Conception rate CHE CF=Interval from Calving to First Service (ICF), days T2=CR NR=Non Return Rate after 56 Days (NRR), % T3=C1 T4=C2 NR=Non Return Rate after 56 Days (NRR), % T1=HC CR=Heifers' Conception rate (pregnant or not after 3 months) CZE T3=C1 CR=Cows' Conception rate (pregnant or not after 3 months) T4=C2 CR=Cows' Conception rate (pregnant or not after 3 months) AUT/DEU T1=HC NR=Heifers' Non Return Rate after 56 days T2=CY CF=Interval from calving to first insemination cows (days) NR=Cows' Non Return Rate after 56 days T3=C1 T4=C2 FL=Interval from first to last insemination cows (days) T5=IT DO=Days open (days) T1=HC NR=Heifers' Non Return Rate after 56 days DFS T2=CY CF=Interval from calving to first insemination cows (days) T3=C1 NR=Cows' Non Return Rate after 56 days T4=C2 FL=Interval from first to last insemination cows (days) T5=IT DO=Days open (days) ESP T2=CY DO=Days open T4=C2 DO=Days open T5=IT DO=Days open FRA T1=HC CR=Heifers' Conception rate (binary trait) for maiden heifers T2=CY Interval between calving and first AI T3=C1 CR=Cows' Conception rate (binary trait) for cows T4=C2 FL=Interval from first to last insemination cows (days) T2=CY CI=days between 1st and 2nd calvings GBR T3=C1 NR=1st lactation non return at 56 days T4=C2 CI=days between 1st and 2nd calvings T5=IT CI=days between 1st and 2nd calvings IRL T2=CY CI=Calving interval T4=C2 CI=Calving interval T5=IT CI=Calving interval ISR T3=C1 CR=Inverse of the number of insemination to conception (%) T4=C2 CR=Inverse of the number of insemination to conception (%) ITA T1=HC NR= non-return rate 56 days (heifers) T2=CY CF=Days to first service T3=C1 NR=Non-return rate at 56 days (%) T4=C2 FL=Interval from first to last insemination cows (days) T5=IT DO=days open (days) ITA(BSW) T2=CY CF=Interval calving to first insemination T4=C2 Days Open T5=IT CI=Calving interval T1=HC CR=Heifers' Conception rate NLD T2=CY CF=Interval calving to first insemination (days) CR=Cows' Conception rate (binary trait) for cows T3=C1 T4=C2 FL=Interval from first to last insemination cows (days) T5=IT CI=Calving Interval (days) T1=HC NR=NR=Non-return rate 56 days (heifers) NOR CF=Interval calving to first insemination (days) T2=CY NR=NR=Non-return rate 56 days (cows) T3=C1 T4=C2 CI=Calving Interval (days) T5=IT CI=Calving Interval (days) T2=CY PM=Lactating cow's ability to start cycling NZL T4=C2 PC=Lactating cow's ability to conceive (CR42) T5=IT PC=Lactating cow's ability to conceive (CR42)
- POL T1=HC CR=Conception rate for heifers

	T2=CR T3=C1 T4=IT T5=IT	Interval from calving to first insemination CR=Conception rate for cows Days open Days open
USA	T1=HC T2=CY T3=C1 T4=C2 T5=IT	CR=Conception rate (heifer) CF=Interval from calving to first insemination CR=Conception rate (cow) DP=Daughter Pregnancy Rate DP=Daughter Pregnancy Rate
ZAF	T4=IT T5=IT	CI=Calving Interval CI=Calving Interval
JPN	T1=HC T2=CY T3=C1 T4=C2 T5=IT	CR=Heifers'Conception rate DO=Days open CR=Cows'Conception rate DO=Days open DO=Days open

CHANGES IN NATIONAL PROCEDURES

\_\_\_\_\_

Changes in the national genetic evaluation of fertility traits are as follows:

CAN	(HOL)	Base change
DFS	(HOL)	Change in status of bulls
FRA	(HOL)	Base change
ITA	(HOL)	Cut off one year of data and base change
NLD	(HOL)	INT: Now send Days open instead of Calving Interval. Conception rate heifers and cows:
		added the used sire to the model and done some minor data edits.
ESP	(HOL)	Exclusion from national genomic evaluation of candidates and culled bulls older than 2 years old
		Reduction in reliability due to reduction of parent average's reliability
DEU	(HOL)	Base change
GBR	(HOL)	Base change
BEL	(HOL)	Base change

INTERBULL CHANGES COMPARED TO THE DECEMBER ROUTINE RUN

-----

No changes in Interbull procedures

DATA AND METHOD OF ANALYSIS

Thirteen Holstein populations sent GEBV data for up to 38 traits, while classical EBVs for the same traits were used in the analyses. Young bull GEBVs from the GEBV providers have been converted to the scales of all countries participating in classical MACE. A bull will get a MACE EBV or a GMACE EBV but not both. From those thirteen countries, National GEBVs of bulls less than seven years of age and with no classical MACE proofs were included for the breeding value prediction with a further requirement of either a MACE-PA or a GMACE-PA (for young genomic bulls with young genomic sires) being available.

The parameter-space approach is used for the GMACE genetic evaluations (Sullivan, 2016)

SCIENTIFIC LITERATURE

The international genetic evaluation procedure is based on international work described in the following scientific publications:

\_\_\_\_\_

Sullivan, P.G. 2016. Defining a Parameter Space for GMACE. Interbull Bulletin 50, p 85-93.

VanRaden, P.M. and Sullivan, P.G. 2010. International genomic evaluation methods for dairy cattle. Gen. Sel. Evol. 42:7

Sullivan, P.G. and Jakobsen, J.H. 2012. Robust GMACE for young bulls methodology. Interbull Bulletin 45, Article 1.

Sullivan, P.G. 2012a. GMACE reliability approximation. Report to the GMACE working group of Interbull. GMACE\_rels 2013 Sullivan, P.G. 2012b. GMACE variance estimation. Report to the GMACE working group of Interbull. GMACE\_vce 2013 Sullivan, P.G. 2012c. GMACE Weighting Factors. Report to the GMACE working group of Interbull. GMACE\_gedcs 2013 Jakobsen, J.H. and Sullivan, P.G. 2013. Trait specific computation of shared reference population. Reference sharing Nov 2013

NEXT ROUTINE INTERNATIONAL EVALUATION

Dates for next routine run can be found on http://www.interbull.org/ib/servicecalendar

NEXT TEST INTERNATIONAL EVALUATION

Dates for next routine run can be found on http://www.interbull.org/ib/servicecalendar

## PUBLICATION OF INTERBULL ROUTINE RUN

Results were distributed by the Interbull Centre to designated representatives in each country. The international evaluation file comprised international proofs expressed on the base and unit of each country included in the analysis. Such records readily provide more information on bull performance in various countries, thereby minimising the need to resort to conversions.

At the same time, all recipients of Interbull results are expected to honour the agreed code of practice, decided by the Interbull Steering Committee, and only publish international evaluations on their own country scale. Evaluations expressed on another country scale are confidential and may only be used internally for research and review purposes.

Table 1. National evaluation dates in GMACE run April 2021

Country	Date
BEL	20201201
CAN	20210401
DEU	20210407
DFS	20210302
ESP	20210319
GBR	20210309
ITA	20210311
NLD	20210401
POL	20210407
FRA	20210407
========	

Table 2.

Number	of	bulls	in	reference	population	for	hco				

CAN 34993.0 DEU 7295.0 39438.0 DFS 4384.0 34498.0 35385.0 FRA 3801.0 32252.0 31713.0 33843.0 POL 4266.0 29715.0 29518.0 27811.0 31554.0 NLD 3870.0 33665.0 33116.0 31747.0 28925.0 34785.0 ITA 28949.0 6367.0 3498.0 3009.0 3319.0 3074.0 29685.0

Number of bulls in reference population for crc

\_\_\_\_\_

 BEL
 1484.0

 CAN
 728.0
 34897.0

 DEU
 717.0
 7579.0
 41911.0

 DFS
 628.0
 4525.0
 36772.0
 37710.0

 ESP
 692.0
 4948.0
 37551.0
 36754.0
 38362.0

 GBR
 669.0
 31491.0
 7815.0
 4734.0
 5155.0
 33746.0

 ITA
 717.0
 31463.0
 6631.0
 3632.0
 4082.0
 30341.0
 32237.0

 NLD
 738.0
 4025.0
 35979.0
 35468.0
 35973.0
 4297.0
 3229.0
 37764.0

 POL
 846.0
 4258.0
 32057.0
 31909.0
 32251.0
 4194.0
 3382.0
 31340.0
 33336.0

 FRA
 700.0
 3996.0
 34347.0
 33822.0
 34395.0
 4061.0
 3193.0
 33871.0
 29960.0
 36046.0

Number of bulls in reference population for cc1

CAN 37943.0 DEU 7490.0 39819.0 DFS 4419.0 34675.0 35525.0 FRA 3900.0 32469.0 31888.0 34123.0 GBR 31498.0 7713.0 4618.0 3963.0 32951.0 ITA 31459.0 6544.0 3541.0 3108.0 30279.0 32204.0 NLD 3940.0 33866.0 33283.0 31921.0 4118.0 3140.0 35075.0 POL 4297.0 29940.0 29728.0 28041.0 4168.0 3354.0 29160.0 31743.0

Number of bulls in reference population for cc2

 BEL
 1677.0

 CAN
 751.0
 40457.0

 DEU
 722.0
 7661.0
 42100.0

 DFS
 630.0
 4583.0
 36890.0
 37834.0

 ESP
 695.0
 5018.0
 37676.0
 38508.0

 GBR
 685.0
 33935.0
 7883.0
 4769.0
 5207.0
 36235.0

 ITA
 731.0
 33195.0
 6660.0
 3645.0
 4100.0
 32043.0
 33969.0

 NLD
 743.0
 4107.0
 36085.0
 35567.0
 36084.0
 4333.0
 3262.0
 38075.0

 POL
 1017.0
 4451.0
 32168.0
 32019.0
 32364.0
 4307.0
 3462.0
 31457.0
 34055.0

Number of bulls in reference population for int

 BEL
 1341.0

 CAN
 578.0
 38562.0

 DEU
 717.0
 7571.0
 41947.0

 DFS
 628.0
 4541.0
 36808.0
 37745.0

 ESP
 692.0
 4969.0
 37588.0
 36804.0
 38412.0

 GBR
 669.0
 33766.0
 7846.0
 4746.0
 5180.0
 36050.0

 ITA
 571.0
 31777.0
 6613.0
 3635.0
 4084.0
 31955.0
 32537.0

 NLD
 738.0
 4043.0
 36009.0
 35495.0
 36003.0
 4311.0
 3248.0
 37933.0

 POL
 767.0
 3979.0
 32085.0
 31940.0
 32280.0
 4200.0
 3119.0
 31365.0
 33077.0