

Introduction

The latest routine international evaluation for SNP Training for clinical mastitis took place as scheduled at the Interbull Centre. Data from six (6) countries were included in this evaluation.

International genetic evaluations for SNP Training for clinical mastitis of bulls from Canada, France, Germany, Netherlands, Switzerland, and the United States of America were computed. Brown Swiss, Holstein and Jersey breed data were included in this evaluation.

Changes in national procedures

Changes in the national genetic evaluation of SNP Training for clinical mastitis are as follows:

No major changes from participating countries

INTERBULL CHANGES COMPARED TO THE PREVIOUS ROUTINE RUN

Post-processing Windows:

According to the decision taken by ITC in Orlando (2015) to review the post-processing windows every 5 years, during the 2020 the relative working group has been re-activated and new windows have been identified.

As before, the upper bounds have been set to 0.99 as these were judged to have very little effect on evaluations while the lower values have been reduced to the 10th percentile. This reduction would provide post-processed correlations to be closer to the real estimated ones. Over the past five years, in fact, the previous adopted lower value (25th percentile) had been found too high causing estimated and post-processed correlations to differ significantly from each other. The new lower values have been applied to all breeds and traits.

The weight assigned to the magnitude of the changes tested by each country has also been revised. The new weight will allow post-processed correlations to take more in consideration the value of the new estimated ones even when no changes are applied by the countries.

The new weights are as follows:

No changes :: 2
Small changes:: 1
Big changes :: 0

More information can be read on https://interbull.org/ib/rg_procedure

DATA AND METHOD OF ANALYSIS

Data were national genetic evaluations of AI sampled bulls with at least 10 daughters or 10 EDC (for clinical mastitis and maternal calving traits at least 50 daughters or 50 EDC, and for direct calving traits at least 50 calvings or 50 EDC) in at least 10 herds. Table 1 presents the amount of data included in this Interbull evaluation for all breeds.

National proofs were first de-regressed within country and then analysed jointly with a linear model including the effects of evaluation country, genetic group of bull and bull merit. Heritability estimates used in both the de-regression and international evaluation were as in each country's national evaluation.

Table 2 presents the date of evaluation as supplied by each country

Estimated genetic parameters and sire standard deviations are shown in APPENDIX I and the corresponding number of common bulls are listed in APPENDIX II.

SCIENTIFIC LITERATURE

The international genetic evaluation procedure is based on international work described in the following scientific publications:

International genetic evaluation computation:
 Schaeffer. 1994. J. Dairy Sci. 77:2671-2678
 Klei, 1998. Interbull Bulletin 17:3-7

Verification and Genetic trend validation:
 Klei et al., 2002. Interbull Bulletin 29:178-182.
 Boichard et al., 1995. J. Dairy Sci. 78:431-437

Weighting factors:
 Fikse and Banos, 2001. J. Dairy Sci. 84:1759-1767

De-regression:
 Sigurdsson and G. Banos. 1995. Acta Agric. Scand. 45:207-219
 Jairath et al. 1998. J. Dairy Sci. Vol. 81:550-562

Genetic parameter estimation:
 Klei and Weigel, 1998, Interbull Bulletin 17:8-14
 Sullivan, 1999. Interbull Bulletin 22:146-148

Post-processing of estimated genetic correlations:
 Mark et al., 2003, Interbull Bulletin 30:126-135
 Jorjani et al., 2003. J. Dairy Sci. 86:677-679
 <https://wiki.interbull.org/public/rG%20procedure?action=print>

Time edits
 Weigel and Banos. 1997. J. Dairy Sci. 80:3425-3430

International reliability estimation
 Harris and Johnson. 1998. Interbull Bulletin 17:31-36

NEXT ROUTINE INTERNATIONAL EVALUATION

Dates for the next routine evaluation can be found on
<http://www.interbull.org/ib/servicecalendar>.

NEXT TEST INTERNATIONAL EVALUATION

Dates for the next test run can be found on
<http://www.interbull.org/ib/servicecalendar>.

PUBLICATION OF INTERBULL ROUTINE RUN

Results were distributed by the Interbull Centre to designated representatives in each country. The international evaluation file comprised international proofs expressed on the base and unit of each country included in the analysis. Such records readily provide more information on bull performance in various countries, thereby minimizing the need to resort to conversions.

At the same time, all recipients of Interbull results are expected to honor the agreed code of practice, decided by the Interbull Steering Committee, and only publish international evaluations on their own country scale. Evaluations expressed on another country scale are confidential and may only be used internally for research and review purposes.

PUBLICATION OF INTERBULL TEST RUN

Test evaluation results are meant for review purposes only and should not be published.

^LTable 1. National evaluation data considered in the Interbull evaluation for SNP training for clinical mastitis (August Routine Evaluation 2021). Number of records for clinical mastitis by breed

| Country | BSW | GUE | HOL | JER | RDC | SIM |
|-------------|------|-----|-------|-----|-----|-----|
| AUS | | | | | | |
| BEL | | | | | | |
| CAN | | | 4914 | 238 | | |
| CHE | 664 | | 646 | | | |
| CZE | | | | | | |
| DEA | | | | | | |
| DEU | | | 4397 | | | |
| DFS | | | | | | |
| ESP | | | | | | |
| EST | | | | | | |
| FRA | 378 | | 12377 | | | |
| FRM | | | | | | |
| GBR | | | | | | |
| HUN | | | | | | |
| IRL | | | | | | |
| ISR | | | | | | |
| ITA | | | | | | |
| JPN | | | | | | |
| KOR | | | | | | |
| LTU | | | | | | |
| LVA | | | | | | |
| NLD | | | 2503 | | | |
| NOR | | | | | | |
| NZL | | | | | | |
| POL | | | | | | |
| PRT | | | | | | |
| SVK | | | | | | |
| SVN | | | | | | |
| URY | | | | | | |
| USA | | | 6976 | 751 | | |
| ZAF | | | | | | |
| HRV | | | | | | |
| CAM | | | | | | |
| No. Records | 1042 | | 31813 | 989 | | |
| Pub. Proofs | 985 | 0 | 27104 | 847 | 0 | 0 |

^LAPPENDIX I. Sire standard deviations in diagonal and genetic correlations below diagonal

| BSW | cma | | | | | |
|-----|-------|-------|-------|------|------|------|
| | CHE | FRA | | | | |
| CHE | 11.28 | | | | | |
| FRA | 0.85 | 1.05 | | | | |
| HOL | cma | | | | | |
| | CAN | CHE | DEU | FRA | USA | NLD |
| CAN | 7.76 | | | | | |
| CHE | 0.90 | 10.51 | | | | |
| DEU | 0.89 | 0.94 | 10.03 | | | |
| FRA | 0.92 | 0.95 | 0.92 | 1.20 | | |
| USA | 0.82 | 0.83 | 0.83 | 0.83 | 2.32 | |
| NLD | 0.83 | 0.93 | 0.84 | 0.86 | 0.82 | 5.01 |

```

-----
JER      cma
-----
      CAN      USA
CAN      8.21
USA      0.83      2.52
-----

```

^LAPPENDIX II. Number of common bulls

```

-----
BSW
-----
common bulls below diagonal
common three quarter sib group above diagonal
      CHE      FRA
-----
      CHE      0      68
      FRA      51      0
-----

```

```

-----
GUE
-----
HOL
-----
common bulls below diagonal
common three quarter sib group above diagonal
      CAN      CHE      DEU      FRA      USA      NLD
-----
      CAN      0      235      776      807      1171      330
      CHE      206      0      236      203      214      135
      DEU      598      213      0      895      744      533
      FRA      600      176      574      0      883      484
      USA      1311      186      620      649      0      345
      NLD      310      119      473      357      308      0
-----

```

```

-----
JER
-----
common bulls below diagonal
common three quarter sib group above diagonal
      CAN      USA
-----
      CAN      0      81
      USA      74      0
-----

```

```

-----
RDC
-----
SIM
-----

```